

## Journal Club

**Editor's Note:** These short reviews of recent *JNeurosci* articles, written exclusively by students or postdoctoral fellows, summarize the important findings of the paper and provide additional insight and commentary. If the authors of the highlighted article have written a response to the Journal Club, the response can be found by viewing the Journal Club at [www.jneurosci.org](http://www.jneurosci.org). For more information on the format, review process, and purpose of Journal Club articles, please see <http://jneurosci.org/content/preparing-manuscript#journalclub>.

## The Impact of Non-Neurotropic Influenza Strains on the Brain: A Role for Microglial Priming?

✉ Maria Carolina Barbosa-Silva,<sup>1</sup> ✉ Luís Eduardo Santos,<sup>2</sup> and ✉ Bárbara Rangel<sup>1</sup>

Institutes of <sup>1</sup>Biophysics and <sup>2</sup>Medical Biochemistry, Federal University of Rio de Janeiro, Rio de Janeiro, RJ 21941-902, Brazil

Review of Hosseini et al.

This year marks the 100th anniversary of the great influenza pandemic of 1918. The “Spanish Flu”, as it came to be known, was one of the deadliest and most widespread in modern history, with fatalities estimated in the tens of millions (Kristensson, 2006). In the wake of the pandemic, an intriguing association, which is yet to be fully understood, began to emerge. A likewise global, decade-long epidemic of encephalitis lethargica—a devastating condition involving sleep disorder, lethargy, and parkinsonism—coincided with the flu, and lasted for the 10 following years, leading many to believe they shared the same causative agent (Ludlow et al., 2016). Numerous cases of other central nervous system (CNS) disorders were also reported in flu patients, collectively suggesting that influenza could be affecting the brain, sometimes to leave lasting sequelae (Henry et al., 2010).

Although the often-cited causal link to encephalitis lethargica has grown contro-

versial (Studahl, 2003; Dale et al., 2004), abundant evidence, including those from recent seasonal outbreaks (Britton et al., 2017) and from the 2009 “Swine Flu” pandemic (Mizuguchi, 2013), have since confirmed that neurological complications may arise as a consequence of influenza infections. Indeed, although influenza viruses are notorious for their respiratory symptoms, their most common extra-respiratory complications are encephalopathies and other CNS conditions, such as myelopathy, ataxia, delirium, and seizures, which, when present, usually appear up to 1 week after the first symptoms of influenza begin (Studahl, 2003; Ludlow et al., 2016).

Some influenza strains are considered neurotropic/neurovirulent, having been shown to enter the CNS through several different means, such as infecting microvascular endothelial cells or ascending the olfactory, vagus, or trigeminal nerves. Interestingly, however, a capacity to enter the CNS and infect neurons directly does not seem to determine whether a strain causes neurological complications (Koyuncu et al., 2013). For instance, despite their well documented association with encephalitis, strains of H1N1 influenza involved in the 2009 pandemic are non-neurotropic (Sadasivan et al., 2015; Wiley et al., 2015).

The most likely explanation for this apparent paradox lies in neuroinflammation. Peripheral cytokine release following

an infection may lead to important consequences in the brain, particularly due to indirect activation or priming of microglia, the local effectors of innate immunity (Santos et al., 2016). Jurgens et al. (2012) showed that this mechanism occurs during influenza infection, by describing the ability of a non-neurotropic H1N1 strain (A/PR/8/34) to produce a central inflammatory response, hippocampal alterations, and concurrent cognitive deficits in mice. These authors, however, only measured short-term effects, up to 7 d post-infection (dpi), in a period marked by the cytokine-induced state of cognitive and behavioral impairment known as “sickness behavior”. Whether cytokine release can result in microglial activation after this acute phase of infection, a potential mechanism for long-term influenza-related neuropathology, remained unanswered.

In a recent issue of *The Journal of Neuroscience*, Hosseini et al. (2018) shed light on this subject by looking into the neurological consequences of influenza up to 120 dpi. Using a mouse model, they compared results from three different strains of the virus, one neurotropic (H7N7), and two non-neurotropic (H1N1 and H3N2). As in the paper by Jurgens et al. (2012; an independent group), Hosseini et al. (2018) focus on the hippocampus and hippocampus-related behavior. Rich in cytokine receptors, this region is particularly sensitive to neuroinflammation (Parnet et al., 2002).

After confirming that, in their model,

Received May 29, 2018; revised July 30, 2018; accepted Aug. 2, 2018.

This work was supported by Grants from Conselho Nacional de Desenvolvimento Científico e Tecnológico and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior; M.C.B.-S. and B.R. are graduate students at the Institute of Biophysics of the Federal University of Rio de Janeiro, and L.E.S. is a postdoctoral fellow at the Institute of Biochemistry of the same University.

The authors declare no competing financial interests.

Correspondence should be addressed to Maria Carolina Barbosa-Silva, Institute of Biophysics, Federal University of Rio de Janeiro, Rio de Janeiro, RJ 21941-902, Brazil. E-mail: mcarolinabarbosa@ufrrj.br.

DOI:10.1523/JNEUROSCI.1368-18.2018

Copyright © 2018 the authors 0270-6474/18/387758-03\$15.00/0

the non-neurotropic strains were indeed unable to enter and replicate effectively in the CNS, Hosseini et al. (2018) detailed parameters of neuroinflammation, synaptic plasticity, and behavior, at 30, 60, and 120 dpi. Among the numerous comparisons and outcomes measured, several novel observations stood out.

At 30 dpi, the density of dendritic spines was reduced in the hippocampus of mice infected with either H7N7 or H3N2 strains. This was accompanied by an increase in the number of microglia and astrocytes in the region. Microglial activation levels, estimated by counting the number of primary processes per cell, were also increased, suggesting gliosis and neuroinflammation. Consistently, infection with H7N7 or H3N2 led to deficits in long-term potentiation and impaired performance in a Morris water maze task at this initial time point. The changes Hosseini et al. (2018) found in the brains of H7N7 and H3N2-infected mice at 30 dpi were partially or completely recovered at 60 dpi. At 120 dpi, all measured parameters were at control levels. Notably, in the H1N1-infected group, none of the parameters were significantly different from control values, even at 30 dpi. But although the observed central effects subsided eventually, their first time point of 30 dpi was already enough to set their results outside the sickness behavior window. This can be inferred from the weight loss (which peaked at 8–9 dpi) and subsequent recovery they observe in infected animals, suggesting that the effects reported are indeed chronic.

The lack of effect in the H1N1 group is noteworthy as it suggests that the alterations in hippocampal morphology previously described by Jurgens et al. (2012) do not persist far beyond the acute phase of infection. Still, it should be noted that these results may not be entirely complementary, as they used mice of a different strain and gender, variables already shown to impact the development of influenza infections (Klein et al., 2012; Samet and Tompkins, 2017).

Despite the eventual recovery, the central effects observed by Hosseini et al. (2018) might have long-term consequences, due to innate immune plasticity. Not unlike macrophages, their peripheral counterparts, microglia can react differently when it has been “primed” by a previous stimulus. This is a complex phenomenon, in which intensity, duration, and frequency of an inflammatory event can render microglia overly sensitive or more tolerant toward future stimuli (Wendeln et al., 2018). Priming

of microglia has been shown to occur indirectly, through the release of peripheral cytokines that reach the CNS, and even as a result of atypical insults, such as the mild inflammation that accompanies normal aging (Luo et al., 2010). Priming effects profoundly affect the brain in several pathological contexts (Perry and Holmes, 2014). Thus, even after H1N1 infection, where no chronic effects were found, microglial priming might lead to long-lasting brain dysfunction and/or increased susceptibility to otherwise inconsequential inflammatory stimuli.

Hosseini et al. (2018) went on to show increased levels of hippocampal TNF- $\alpha$  at 10 dpi, in all the influenza-infected groups. This local increase in brain TNF- $\alpha$ , which was similar to results shown by Jurgens et al. (2012), further suggests that microglial priming may take place in this model of influenza infection. TNF- $\alpha$  activates microglia in an autocrine loop (Kuno et al., 2005) and, along with other inflammatory cytokines, could indeed precipitate the lasting changes in gene expression profiles that amount to priming (Perry and Holmes, 2014). Future work should investigate whether such priming occurs, especially given that multiple influenza infections may happen throughout a patient's life.

Whether they reach the brain from the periphery or are locally overproduced by activated microglia, inflammatory mediators are known to interfere with memory function. This may happen through several signaling mechanisms, involving, for example, lowering expression of BDNF, an important mediator of synaptic plasticity; activating the IDO enzyme, which ultimately reduces serotonin availability; or directly inhibiting neurogenesis (Santos et al., 2016). These effects are likely to be at play to some extent in the earlier neuroinflammation-induced neurological consequences of influenza infection and could perhaps be a starting point to the more durable deficits now described.

Finally, Hosseini et al. (2018) evaluated whole-genome mRNA data from the hippocampus of H3N2- and H7N7-infected animals and their controls, at 18 and 30 dpi. These results offered insight into possible mechanisms mediating chronic viral effects. Among several interesting hits, two alterations, which apparently do not depend on viral tropism, stand out. BDNF was indeed reduced by infection and may explain, at least partly, both the morphological and behavioral effects observed. Additionally, a robust increase in expression of the dopamine

transporter gene was observed in influenza-infected mice. If this increase indeed translates to altered levels of extracellular dopamine, it will most likely have a meaningful role not only in memory and learning behavior, but also in other types of behavior and related processes, such as neurogenesis (Borta and Höglinger, 2007; Rossato et al., 2009). Moreover, given that monoamine neurotransmitter transporters may be unspecific, and that monoaminergic systems may interact deeply with one another (Daws, 2009), this increase may have consequences to serotonergic and noradrenergic synapses as well.

Even a century after the great pandemic, the threat of another highly virulent strain emerging is always looming. Despite continued efforts to develop vaccines against such a highly mutable virus, influenza remains a leading cause of morbidity and mortality around the world. Because of this, understanding the long-term impacts of infection, particularly in the CNS, is paramount. The report by Hosseini et al. (2018) provides evidence that even common non-neurotropic strains of influenza can cause lasting changes to the hippocampus, interfering with normal behavior after the acute infection is resolved. Whether there is a causal role for peripheral inflammation and microglia in this effect is an interesting possibility that remains to be seen. Further research on the neurological consequences of influenza will not only help us understand the virus, but also how the brain reacts to peripheral inflammation.

## References

- Borta A, Höglinger GU (2007) Dopamine and adult neurogenesis. *J Neurochem* 100: 587–95. [CrossRef Medline](#)
- Britton PN, Blyth CC, Macartney K, Dale RC, Li-Kim-Moy J, Khandaker G, Crawford NW, Marshall H, Clark JE, Elliott EJ, Booy R, Cheng AC, Jones CA (2017) The spectrum and burden of influenza-associated neurological disease in children: combined encephalitis and influenza sentinel site surveillance from Australia, 2013–2015. *Clin Infect Dis* 65:653–660. [CrossRef Medline](#)
- Dale RC, Church AJ, Surtees RA, Lees AJ, Adcock JE, Harding B, Neville BG, Giovannoni G (2004) Encephalitis lethargica syndrome: 20 new cases and evidence of basal ganglia autoimmunity. *Brain* 127:21–33. [CrossRef Medline](#)
- Daws LC (2009) Unfaithful neurotransmitter transporters: focus on serotonin uptake and implications for antidepressant efficacy. *Pharmacol Ther* 121: 89–99. [CrossRef Medline](#)
- Henry J, Smeyne RJ, Jang H, Miller B, Okun MS (2010) Parkinsonism and neurological manifestations of influenza throughout the 20th and 21st centuries. *Parkinsonism Relat Disord* 16:566–571. [CrossRef Medline](#)

- Hosseini S, Wilk E, Michaelsen-Preusse K, Gerhauser I, Baumgärtner W, Geffers R, Schughart K, Korte M (2018) Long-term neuroinflammation induced by influenza A virus infection and the impact on hippocampal neuron morphology and function. *J Neurosci* 38: 3060–3080. [CrossRef Medline](#)
- Jurgens HA, Amacherla K, Johnson RW (2012) Influenza infection induces neuroinflammation, alters hippocampal neuron morphology, and impairs cognition in adult mice. *J Neurosci* 32: 3958–3968. [CrossRef Medline](#)
- Klein SL, Hodgson A, Robinson DP (2012) Mechanisms of sex disparities in influenza pathogenesis. *J Leukoc Biol* 92:67–73. [CrossRef Medline](#)
- Koyuncu OO, Hogue IB, Enquist LW (2013) Virus infections in the nervous system. *Cell Host Microbe* 13:379–393. [CrossRef Medline](#)
- Kristensson K (2006) Avian influenza and the brain: comments on the occasion of resurrection of the spanish flu virus. *Brain Res Bull* 68:406–413. [CrossRef Medline](#)
- Kuno R, Wang J, Kawanokuchi J, Takeuchi H, Mizuno T, Suzumura A (2005) Autocrine activation of microglia by tumor necrosis factor- $\alpha$ . *J Neuroimmunol* 162: 89–96. [CrossRef Medline](#)
- Ludlow M, Kortekaas J, Herden C, Hoffmann B, Tappe D, Trebst C, Griffin DE, Brindle HE, Solomon T, Brown AS, van Riel D, Wolthers KC, Pajkrt D, Wohlsein P, Martina BEE, Baumgärtner W, Verjans GM, Osterhaus ADME (2016) Neurotropic virus infections as the cause of immediate and delayed neuropathology. *Acta Neuropathol* 131:159–184. [CrossRef Medline](#)
- Luo XG, Ding JQ, Chen SD (2010) Microglia in the aging brain: relevance to neurodegeneration. *Mol Neurodegener* 5:12. [CrossRef Medline](#)
- Mizuguchi M (2013) Influenza encephalopathy and related neuropsychiatric syndromes. *Influenza Other Respir Viruses* 7:67–71. [CrossRef Medline](#)
- Parinet P, Kelley KW, Bluthé RM, Dantzer R (2002) Expression and regulation of interleukin-1 receptors in the brain. Role in cytokines-induced sickness behavior. *J Neuroimmunol* 125:5–14. [CrossRef Medline](#)
- Perry VH, Holmes C (2014) Microglial priming in neurodegenerative disease. *Nat Rev Neurol* 10: 217–24. [CrossRef Medline](#)
- Rossato JI, Bevilacqua LR, Izquierdo I, Medina JH, Cammarota M (2009) Dopamine controls persistence of long-term memory storage. *Science* 325:1017–1020. [CrossRef Medline](#)
- Sadasivan S, Zanin M, O'Brien K, Schultz-Cherry S, Smeyne RJ (2015) Induction of microglia activation after infection with the non-neurotropic A/CA/04/2009 H1N1 influenza virus. *PLoS One* 10: e0124047. [CrossRef Medline](#)
- Samet SJ, Tompkins SM (2017) Influenza pathogenesis in genetically defined resistant and susceptible murine strains. *Yale J Biol Med* 90: 471–479. [Medline](#)
- Santos LE, Beckman D, Ferreira ST (2016) Microglial dysfunction connects depression and Alzheimer's disease. *Brain Behav Immun* 55: 151–165. [CrossRef Medline](#)
- Studahl M (2003) Influenza virus and CNS manifestations. *J Clin Virol* 28:225–232. [CrossRef Medline](#)
- Wendeln AC, Degenhardt K, Kaurani L, Gertig M, Ulas T, Jain G, Wagner J, Häslér LM, Wild K, Skodras A, Blank T, Staszewski O, Datta M, Centeno TP, Capece V, Islam MR, Kerimoglu C, Staufienbiel M, Schultze JL, Beyer M, et al. (2018) Innate immune memory in the brain shapes neurological disease hallmarks. *Nature* 556:332–338. [CrossRef Medline](#)
- Wiley CA, Bhardwaj N, Ross TM, Bissel SJ (2015) Emerging infections of CNS: avian influenza A virus, rift valley fever virus and human parechovirus. *Brain Pathol* 25:634–650. [CrossRef Medline](#)